

Chemotherapy Orders ETHYOL (amifostine) and CISPLATIN

Date chemotherapy to be given: _____

Weight: _____	<input type="checkbox"/> lbs.	<input type="checkbox"/> kg	Height: _____	<input type="checkbox"/> inches	<input type="checkbox"/> cm
	<input type="checkbox"/> actual	<input type="checkbox"/> ideal			
BSA used for calculations: _____m ²					

For cycle 2 and subsequent cycles , are doses identical to previous cycle? <input type="checkbox"/> Yes <input type="checkbox"/> No If no, indicate reason for adjustment: _____
--

Treat only if: WBC ≥ _____ ANC ≥ _____ Platelets ≥ _____ Other _____
--

<p>Premedicate with:</p> <p>Anzemet _____ mg IVP or Zofran _____ mg IVPB</p> <p>Decadron _____ mg slow IVP or in _____ cc NS or _____ over 30 minutes</p> <p>Other _____</p> <p>Ethyol (amifostine) _____ mg/m² (<i>usual dose 740 mg/m²</i>) x _____ m² = _____ mg in _____ cc NS or _____ over 2-5 minutes then</p> <p>Phenergan _____ mg IVP then</p> <p>250ml NS or _____ over 15 minutes immediately after Ethyol</p> <p>Keep patient supine during and 15 minutes after Ethyol Check BP every minute during Ethyol. Interrupt infusion if systolic BP drops 20 mmHg from baseline during Ethyol infusion and infuse NS at 500cc/hr until BP returns to baseline. If after 5 minutes and patient is asymptomatic, resume infusion. If after 5 minutes BP does not return to baseline, infusion should be terminated.</p> <p>Within 30 minutes after Ethyol begin Cisplatin _____ mg/m² (<i>usual dose 20-40 mg/m²</i>) x _____ m² = _____ mg in _____ cc NS or _____ over _____ minutes.</p> <p>_____(<i>check if desired</i>) Add 25 gram mannitol and 2 gram magnesium sulfate to same bag as Cisplatin.</p> <p>After Cisplatin is finished run _____ cc NS or _____ over 1 hour then resume prior IV rate.</p>

Physician Signature: _____ **Date:** _____



CHEMOTHERAPY ORDERS ETHYOL (amifostine) and CISPLATIN